

Sarah Clark

## ORIGINAL ARTICLE

## Weight Lifting in Women with Breast-Cancer-Related Lymphedema

Kathryn H. Schmitz, Ph.D., M.P.H., Rehana L. Ahmed, M.D., Ph.D.,  
 Andrea Troxel, Sc.D., Andrea Cheville, M.D., Rebecca Smith, M.D.,  
 Lorita Lewis-Grant, M.P.H., M.S.W., Cathy J. Bryan, M.Ed.,  
 Catherine T. Williams-Smith, B.S., and Quincy P. Greene

## ABSTRACT

## BACKGROUND

From the Center for Clinical Epidemiology and Biostatistics, University of Pennsylvania School of Medicine and Abramson Cancer Center, Philadelphia (K.H.S., A.T., R.S., L.L.-G., C.J.B., C.T.W.-S., Q.P.G.); the Department of Dermatology, University of Minnesota Medical School, Minneapolis (R.L.A.); and the Department of Physical Medicine and Rehabilitation, Mayo Clinic, Rochester, MN (A.C.). Address reprint requests to Dr. Schmitz at the Department of Biostatistics and Epidemiology, University of Pennsylvania, 423 Guardian Dr., 903 Blockley Hall, Philadelphia, PA 19104-6021, or at schmitz@mail.med.upenn.edu.

Weight lifting has generally been proscribed for women with breast-cancer-related lymphedema, preventing them from obtaining the well-established health benefits of weight lifting, including increases in bone density.

## METHODS

We performed a randomized, controlled trial of twice-weekly progressive weight lifting involving 141 breast-cancer survivors with stable lymphedema of the arm. The primary outcome was the change in arm and hand swelling at 1 year, as measured through displaced water volume of the affected and unaffected limbs. Secondary outcomes included the incidence of exacerbations of lymphedema, number and severity of lymphedema symptoms, and muscle strength. Participants were required to wear a well-fitted compression garment while weight lifting.

## RESULTS

The proportion of women who had an increase of 5% or more in limb swelling was similar in the weight-lifting group (11%) and the control group (12%) (cumulative incidence ratio, 1.00; 95% confidence interval, 0.88 to 1.13). As compared with the control group, the weight-lifting group had greater improvements in self-reported severity of lymphedema symptoms ( $P=0.03$ ) and upper- and lower-body strength ( $P<0.001$  for both comparisons) and a lower incidence of lymphedema exacerbations as assessed by a certified lymphedema specialist (14% vs. 29%,  $P=0.04$ ). There were no serious adverse events related to the intervention.

## CONCLUSIONS

In breast-cancer survivors with lymphedema, slowly progressive weight lifting had no significant effect on limb swelling and resulted in a decreased incidence of exacerbations of lymphedema, reduced symptoms, and increased strength. (ClinicalTrials.gov number, NCT00194363.)

N Engl J Med 2009;361:664-73.

Copyright © 2009 Massachusetts Medical Society.

## WEIGHT LIFTING IN WOMEN WITH BREAST-CANCER-RELATED LYMPHEDEMA

**T**HERE ARE MORE THAN 2.4 MILLION SURVIVORS of breast cancer in the United States.<sup>1</sup> Approximately 184,000 women are diagnosed with breast cancer each year in the United States, and 90% of these women will live at least 5 years.<sup>2</sup> Improvements in immediate treatment outcomes have led to an increased focus on morbidity among survivors. Lymphedema ranks high among the serious concerns of survivors, as it is chronic, progressive, and incurable.<sup>3</sup> Lymphedema causes limb swelling and discomfort, considerably impairing arm function.<sup>4</sup> The incidence of lymphedema after breast-cancer surgery varies across studies from 6 to 70%, depending on the criteria used for diagnosis and the follow-up interval.<sup>5-7</sup> Sentinel-lymph-node biopsy is associated with a lower risk of lymphedema than axillary dissection,<sup>8</sup> but one study reported a 17% incidence after sentinel-lymph-node biopsy alone.<sup>9</sup>

Breast-cancer survivors with lymphedema may limit the use of their affected arm out of fear and on the basis of guidance from commonly accessed cancer-information Web sites,<sup>10,11</sup> which suggest that risk of lymphedema is decreased by avoiding lifting children, heavy bags, or other objects with the affected arm. Although this advice is intended to prevent harm, adherence to these precautions may limit physical recovery after breast-cancer surgery, alter activities, and adversely affect employment. Conversely, a program of controlled exercise through weight lifting may increase the physical-work capacity of the affected arm, thereby protecting it from injury sustained during common daily activities. Weight lifting offers additional benefits particularly relevant to breast-cancer survivors, including control of body fat<sup>12,13</sup> and improved functional outcomes and bone health.<sup>14-16</sup> We performed a 1-year randomized, controlled trial involving breast-cancer survivors with lymphedema to assess the effects of controlled weight lifting.

---

## METHODS

---

### PATIENTS

A total of 141 women with a history of breast cancer and current lymphedema were recruited from October 2005 through March 2007; follow-up was completed by August 2008. Recruitment methods included letters sent by state cancer registries, advertisements and interviews, and flyers.

Eligible women had a history of unilateral non-metastatic breast cancer 1 to 15 years before study entry and a body-mass index (BMI, the weight in kilograms divided by the square of the height in meters) of 50 or less, were not actively trying to lose weight, and had no current evidence of cancer, no medical conditions that would limit exercise, no history of weight lifting during the previous year, at least one lymph node removed, and a clinical diagnosis of stable breast-cancer-related lymphedema. Lymphedema was defined as a difference in the volume or circumference between the affected and unaffected limb of 10% or more or, according to Common Toxicity Criteria,<sup>17</sup> arm swelling, obscuration of the anatomical architecture of the arm, or pitting edema. If a woman reported having lymphedema but it was not evident at study entry, she was required to provide written documentation of a previous clinical diagnosis of lymphedema and treatment from a certified lymphedema therapist.<sup>18</sup> Stable lymphedema was defined as the absence in the past 3 months of therapist-delivered treatment, more than one arm infection requiring antibiotics, change in ability to perform activities of daily living, and verified changes in arm swelling of more than 10%.


Figure 1 shows the enrollment, randomization, and follow-up of the study participants. Women were assigned to either of two equal-size groups through a computerized process called minimization,<sup>19,20</sup> in a manner that was unpredictable and was concealed from the study staff who determined eligibility. This approach balanced important potential confounders at baseline: age (<54 years vs. ≥54 years), difference in the volume between the affected and unaffected limbs (<10% vs. 10 to 20% vs. >20%), number of lymph nodes removed (<6 vs. ≥6), obesity (BMI <30 vs. ≥30), months since diagnosis (<60 vs. ≥60), and history of radiation treatment (yes vs. no).

The Institutional Review Board of the University of Pennsylvania approved the protocol. Before participating, women provided written informed consent and written clearance from a physician.

### MEASUREMENTS

Measurements were obtained for all participants at baseline and at 12 months by trained staff who were unaware of the study-group assignments, using standardized methods. Limb volume was measured by submerging the arm and hand in

## The NEW ENGLAND JOURNAL of MEDICINE



water and measuring the displaced water volume.<sup>21,22</sup> All participants were also evaluated by a certified lymphedema therapist<sup>18</sup> using the Common Toxicity Criteria, which assess the tone, texture, and anatomical architecture of the arm tissue, in addition to swelling.<sup>27</sup> Participants com-

pleted a validated survey assessing the presence and severity of 14 lymphedema-related limb symptoms (including swelling, leathery skin texture, heaviness, pain, pitting, and difficulty writing).<sup>23</sup> Lymphedema exacerbations were ascertained by certified lymphedema specialists<sup>18</sup> who were

## WEIGHT LIFTING IN WOMEN WITH BREAST-CANCER-RELATED LYMPHEDEMA

unaware of the study-group assignments, using a standardized evaluation. The participant was deemed to have an exacerbation if there was an increase in the volume of the affected limb of 5% or more, accompanied by an increase of 5% or more in the difference in the volume or circumference between the affected and unaffected limbs and by indications of sustained tissue changes such as fibrosis (sponginess, pitting, or hard, nonpitting fibrosis), altered skin color, or alteration of activities of daily living over the previous week because of symptoms (e.g., heaviness, inability to grip, tiredness, or achiness). Participants were evaluated for possible exacerbation if they reported a change in symptoms lasting 1 week or longer or if interim measurements at 3 or 6 months, by staff who were unaware of the study-group assignments, indicated an increase in the volume of the affected limb and an interlimb difference of 5% or more.

For each participant, the maximum amount of weight that could be lifted once was assessed for the bench press and leg press at baseline and 12 months; these tests are considered safe for most populations if properly supervised.<sup>24-26</sup> The initial weight attempted for this test was based on the participant's rating of the difficulty of a warm-up set of four to six repetitions (40 lb [18.1 kg] for leg press, 5 lb [2.3 kg] for bench press), performed after stretching and familiarization with the equipment. Resistance was added until the participant rated the difficulty as maximal and refused to try to lift more, was clearly unable to lift more with proper biomechanics, or reported a symptom that required stopping. Trained study staff encouraged participants according to a standardized script. Compliance with the weight-lifting intervention was evaluated by means of attendance logs completed by fitness trainers.

Demographic characteristics were self-reported by patients at baseline. The cancer stage was obtained for each patient from state registries. Treatment history was self-reported, except for the number of lymph nodes removed, which was derived from pathology reports. Measurements assessed at baseline and at 12 months included weight and height, whole-body dual-energy x-ray absorptiometry (Hologic Discovery), adherence to common lymphedema self-care activities (e.g., compression, massage, bandaging), physical activity outside of weight lifting (according to the

International Physical Activity Questionnaire)<sup>27</sup>; and diet (according to the Diet History Questionnaire).<sup>28,29</sup>

## WEIGHT-LIFTING PROGRAM

Participants assigned to the weight-lifting group received a 1-year membership at a community fitness center (e.g., a YMCA) near their home. For the first 13 weeks, women were instructed, in small groups in a 90-minute session, twice weekly. Certified fitness professionals employed by the fitness centers led these sessions, which included stretching, cardiovascular warm-up, abdominal and back exercises, and weight-lifting exercises. Upper-body exercises included seated row, chest press, lateral or front raises, bicep curls, and tricep pushdowns. Lower-body exercises included leg press, back extension, leg extension, and leg curl. Weight-lifting exercises were introduced with little-to-no resistance. One to three new exercises were taught per session.

During the first 5 weeks, participants increased their number of sets of each exercise per session from two to three, with 10 repetitions per set. If no changes in symptoms were noted for a particular exercise after two sessions at a given weight, the resistance was increased by the smallest possible increment. If fatigue prevented the completion of a third set of 10 repetitions of a given exercise with proper biomechanical form, resistance for that exercise would remain the same at the next session. After two sessions at which three sets of 10 repetitions could be performed with proper form at a given level of resistance, without changes in arm and hand symptoms, the trainer guided the participant to increase the resistance by the smallest possible increment at the next session. No upper limit was placed on the weight to which women could progress in any exercise. During lymphedema exacerbations, women continued all exercises except the upper-body exercises, which were resumed only after approval of their lymphedema therapist, with resistance reset to the lowest possible level and then increased again as described above.

After the first 13 weeks, participants continued twice-weekly unsupervised exercise for 39 additional weeks. Throughout the study, fitness trainers telephoned women who missed more than one session per week. Participants in the control group were asked not to change their exercise level during study participation and were

Table 2. Strength, Anthropometric, Diet, and Physical-Activity Data at Baseline and 12 Months, According to Study Group.\*

Variable	Baseline			12 Mo			% Change between Baseline and 12 Mo			P Value
	Weight Lifting no. with data	Control mean ±SD	P Value	Weight Lifting no. with data	Control mean ±SD	P Value	Weight Lifting no. with data	Control mean ±SD	P Value	
Bench press, maximum (lb)	70	43±16	0.21	56	53±18	<0.001	56	29.4±36.5	<0.001	
Leg press, maximum (lb)	70	182±64	0.09	59	232±66	<0.001	59	32.5±33.2	<0.001	
Weight (kg)	71	82.4±16.6	0.18	65	80.0±15.6	0.66	65	-1.0±3.7	0.47	
BMI†	71	31.0±6.2	0.21	65	30.0±5.7	0.70	65	-1.0±3.7	0.42	
Body fat (%)	71	40.1±5.7	0.42	65	39.6±6.3	0.90	65	-0.3±5.7	0.19	
Fat mass (kg)	71	33.3±10.3	0.27	65	32.0±10.1	0.87	65	-1.2±8.0	0.27	
Lean mass (kg)	71	50.3±7.7	0.29	65	49.1±7.3	0.63	65	-1.0±3.4	0.67	
Diet (kcal per day)	70	1835±1319	0.36	60	1532±679	0.08	60	-1.1±39.2	0.36	
Physical-activity level (MET- min/wk)‡	54	1954±2423	0.49	52	2397±2548	0.20	48	2.8±16.8	0.26	

\* To convert bench and leg press values to kilograms, multiply by 0.45359. MET denotes metabolic equivalent.

† The body-mass index (BMI) is the weight in kilograms divided by the square of the height in meters.

‡ The percent changes over the 12-month study period were calculated from natural-log-transformed physical-activity levels to improve the normality of the distribution.

(Table 2). Changes in diet, physical activity and anthropometric measures over the 12-month period were not significantly different between the two groups.

There was no significant difference between the two groups in the proportion of women who had a change in limb swelling of 5% or more (Table 3). This result did not materially change when the analysis was repeated without any imputed data from participants who were lost to follow-up (results not shown). Among the 130 women who had no second primary or recurrent cancers and were not lost to follow-up, 23 in the control group and 20 in the weight-lifting group were evaluated for exacerbation. Of these, 19 and 9 participants, respectively, were found to have had an exacerbation (83% vs. 45%). The total number of treatment sessions for exacerbation was 195 in the control group, as compared with 77 in the weight-lifting group. The number and severity of symptoms reported decreased more in the weight-lifting group than in the control group. No significant differences were noted between the two study groups with regard to self-reported adherence to prescribed lymphedema self-care therapies (results not shown). Post hoc analyses that excluded participants with grade 0 lymphedema yielded results similar to those reported in Table 3 (results not shown). Adjustment for baseline variables (cancer stage, number of nodes removed, race, physical activity, diet, and body-mass index) did not materially alter these results. There were no serious adverse events related to the intervention.

DISCUSSION

Contrary to common guidelines to avoid lifting with the affected limb, we found that weight lifting did not significantly affect the severity of breast cancer-associated lymphedema (as assessed by the primary outcome, an absolute increase of ≥5 percentage points in the interlimb volume difference). In addition, weight lifting reduced the number and severity of arm and hand symptoms, increased muscular strength, and reduced the incidence of lymphedema exacerbations as assessed by a lymphedema specialist.

Several previous studies, including a case series<sup>33</sup> and small randomized, controlled trials,<sup>34-37</sup> have also suggested that weight lifting is safe for breast-cancer survivors with lymphedema. The

## WEIGHT LIFTING IN WOMEN WITH BREAST-CANCER-RELATED LYMPHEDEMA

Table 3. Lymphedema Outcomes at 12 Months, According to Study Group.\*

Variable	Weight Lifting		Control		Cumulative Incidence Ratio or Mean Difference (95% CI)†	P Value‡
	no. of patients with data	value	no. of patients with data	value		
Change in interlimb volume difference						
≥5% increase — no. (%)	70	8 (11)	69	8 (12)	1.00 (0.88 to 1.13)	1.00
≥5% decrease — no. (%)	70	13 (19)	69	15 (22)	0.96 (0.81 to 1.14)	0.68
Mean interlimb volume discrepancy between baseline and 12 mo (percentage points)	70	-0.69±5.87	69	-0.98±7.31	-0.29 (-1.94 to 2.51)	0.80
Exacerbation — no. (%)	65	9 (14)	65	19 (29)	0.47 (0.23 to 0.97)	0.04
Change in no. of symptoms reported between baseline and 12 mo§	70	-1.81±2.16	69	-1.17±1.94	-0.63 (-1.32 to 0.06)	0.07
Change in severity of symptoms between baseline and 12 mo§	70	-0.51±0.80	69	-0.22±0.71	-0.29 (-0.54 to -0.03)	0.03

\* Plus-minus values are means ±SD.

† The mean difference is given for the weight-lifting group as compared with the control group for the difference in interlimb volume discrepancies (the interarm difference over time) and changes in number and severity of symptoms. The cumulative incidence ratio is given for the weight-lifting group as compared with the control group for differences in percentages.

‡ P values were calculated with the use of Fisher's exact test for between-group comparisons of percentages and the Wilcoxon rank-sum test for between-group comparisons of the difference in interlimb volume discrepancies and changes in number and severity of symptoms.

§ Data were reported by patients regarding 14 symptoms: rings too tight, watch too tight, bracelets too tight, clothing too tight, puffiness, knuckles not visible, veins not visible, skin feels leathery, arm feels tired, pain, pitting, swelling after exercise, difficulty writing, or other. The change in severity of symptoms is the mean of the changes in severity for all 14 symptoms, with the possible severity score for each ranging from 0 (no symptom) to 4 (very severe).

current trial was larger and of longer duration than those previously reported and also differed by testing a weight-lifting protocol with no upper limit on the resistance level to which participants could progress. A strength of this trial is its delivery in community fitness centers, primarily YMCAs, by trainers employed by these fitness centers. We adopted this approach with the goal of dissemination of the weight-lifting program if it proved effective. The ongoing LIVESTRONG at the YMCA program (a collaboration of the YMCA and the Lance Armstrong Foundation) includes the protocol described here as an intervention that can be offered to cancer survivors in YMCAs across the United States. Additional strengths of the present trial are the inclusion of a racially diverse population with a wide range of time since diagnosis (1 to 15 years) and the high rate of follow-up.

There are also potential limitations of the study. Evaluations for exacerbations were not completed by a single therapist, although the six lymphedema therapists assessing exacerbations followed a standardized algorithm for evaluation and had completed the 135-hour course recommended by the National Lymphedema Network.<sup>18</sup>

Therapists were unaware of which patients had been assigned to the weight-lifting group, as specified in the study design, but some participants in this group may have disclosed their recent weight lifting during evaluations for perceived exacerbations. Though the number of women evaluated for exacerbation was approximately equal in the two groups (23 in the control group and 20 in the weight-lifting group), the proportion of evaluated women who were found to have had an exacerbation was higher in the control group. One possible explanation for this observation is that some assessors may have become aware of the study-group assignments, resulting in biased assessments. However, the finding that symptom severity improved more in the weight-lifting group than in the control group supports a benefit of the intervention. An alternative explanation is that participants in the weight-lifting group, concerned about the potential for worsening of lymphedema with weight lifting, were more likely to seek care in the absence of objective evidence of exacerbation.

Although reporting bias cannot be ruled out as a possible explanation for the decrease in confirmed lymphedema exacerbations, several phys-

## WEIGHT LIFTING IN WOMEN WITH BREAST-CANCER-RELATED LYMPHEDEMA

- assessment in the elderly. *J Cardiopulm Rehabil* 1995;15:283-7.
27. Craig CL, Marshall AL, Sjöström M, et al. International Physical Activity Questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003;35:1381-95.
28. Subar AF, Midthune D, Kulidorff M, et al. Evaluation of alternative approaches to assign nutrient values to food groups in food frequency questionnaires. *Am J Epidemiol* 2000;152:279-86.
29. Subar AF, Thompson FB, Kipnis V, et al. Comparative validation of the Block, Willett and National Cancer Institute food frequency questionnaires: the Eating at America's Table Study. *Am J Epidemiol* 2001;154:1089-99.
30. Exercise. Oakland, CA: National Lymphedema Network, 2005. (2007 Update accessed July 20, 2009, at <http://www.lymphnet.org/pdfDocs/nlnextercise.pdf>.)
31. Treatment. Oakland, CA: National Lymphedema Network, 2006. (Accessed July 20, 2009, at <http://www.lymphnet.org/pdfDocs/nlntreatment.pdf>.)
32. Lymphedema risk reduction practices. Oakland, CA: National Lymphedema Network, 2008. (Accessed July 20, 2009, at <http://www.lymphnet.org/pdfDocs/nlntreatment.pdf>.)
33. Harris SR, Niesen-Vertommen SL. Challenging the myth of exercise-induced lymphedema following breast cancer: a series of case reports. *J Surg Oncol* 2000;74:95-9.
34. Ahmed RL, Thomas W, Yee D, Schmitz K. Randomized controlled trial of weight training and lymphedema in breast cancer survivors. *J Clin Oncol* 2006;24:2765-72. [Erratum, *J Clin Oncol* 2006;24:3716.]
35. Johansson K, Tibe K, Weibull A, Newton RC. Low intensity resistance exercise for breast cancer patients with arm lymphedema with or without compression sleeve. *Lymphology* 2005;38:167-80.
36. Courneya KS, Segal RJ, Mackey JR, et al. Effects of aerobic and resistance exercise in breast cancer patients receiving adjuvant chemotherapy: a multicenter randomized controlled trial. *J Clin Oncol* 2007;25:4396-404.
37. McKenzie DC, Kalda AL. Effect of upper extremity exercise on secondary lymphedema in breast cancer patients: a pilot study. *J Clin Oncol* 2003;21:463-6.
38. Mortimer PS. Assessment of peripheral lymph flow before and after clinical intervention. In: *Progress in lymphology*. Amsterdam: Elsevier Science, 1990:215-522.
39. *Idem*. Evaluation of lymphatic function: abnormal lymph drainage in venous disease. *Int Angiol* 1995;14:Suppl 1:32-5.
40. Bourgeois P, Leduc O, Leduc A. Manual of lymphatic drainage: scintigraphic demonstration of its efficacy on colloidal protein resorption. In: *Progress in lymphology*. Amsterdam: Elsevier Science, 1988:551-4.
41. Brennan MJ, Miller LT. Overview of treatment options and review of the current role and use of compression garments, intermittent pumps, and exercise in the management of lymphedema. *Cancer* 1998;83:Suppl:2821-7.

Copyright © 2009 Massachusetts Medical Society.

#### ICMJE SEEKING TWO NEW MEMBER JOURNALS

The International Committee of Medical Journal Editors (ICMJE) is seeking two new member journals to be represented by their editors-in-chief. Information about the ICMJE is available at [www.icmje.org](http://www.icmje.org). The ICMJE anticipates selection of new members by November 1, 2009. Candidate journals should meet the following criteria:

- peer reviewed, general medical journal
- represent geographic areas (Latin America, Asia, Africa) or publication models (open access) not well represented by current ICMJE members
- editor who is knowledgeable about publication ethics
- editor who expects to be in the position for at least 3 years

To apply, editors-in-chief of interested journals should send electronic copies of the following to the ICMJE secretariat (Christine Laine at [claine@acponline.org](mailto:claine@acponline.org)) by September 1:

- brief curriculum vitae
- description of journal, including age, sponsor/publisher, publishing model (subscription model, author pays, open access, etc.), target audience, circulation, number of manuscript submissions per year, description of peer review process used to select material for publication, acceptance rate, bibliographical databases where indexed, Web site address if applicable, and copy of guidelines for authors
- statement on why the journal/editor-in-chief wants to be an ICMJE member (should not exceed 1000 words in length)
- contact information